

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/18120

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A61K 39/395, 38/00; C07K 16/00

US CL : 424/130.1; 514/2; 530/387.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/130.1; 514/2; 530/387.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	US 5,736,137 A (ANDERSON et al) 07 April 1998. See entire document.	1, 2, 7-11, 17, 19-26 ----- 3-6, 12-16, 18, 27-31
Y	MUZAFFAR et al. Immunophenotypic analysis of non-Hodgkin's lymphoma. J. Pak. Med. Assoc. April 1997, Vol. 47, No. 4, pages 108-109. See entire document.	1-31
Y	JUWEID et al. Treatment of non-Hodgkin's lymphoma with radiolabeled murine, chimeric, or humanized LL2, an anti-CD22 monoclonal antibody. Cancer Research. 01 December 1995, Vol. 55 (Suppl.), pages 5899s-5907s. See entire document.	1-31



Further documents are listed in the continuation of Box C.



See patent family annex.

* "A" "E" "L" "O" "P"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance earlier document published on or after the international filing date document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published prior to the international filing date but later than the priority date claimed	*T* "X" "Y" "&"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art document member of the same patent family
--------------------------------------	---	--------------------------	--

Date of the actual completion of the international search

23 NOVEMBER 1999

Date of mailing of the international search report

27 JAN 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MINH-TAM DAVIS

Telephone No. (703) 308-0196



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/18120

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NIELSEN et al. Interferon- α -induced changes in surface antigens in a hairy-cell leukemia (JOK-1), and a Burkitt's lymphoma cell line (Daudi) during <i>in vitro</i> culture. Eur. J. Haematol. 1992, Vol. 48, No. 3, pages 146-151. See entire document.	12-16



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/18120

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST, DIALOG

search terms: anti-CD20 antibody, B-cell lymphoma, non-Hodgkin's lymphoma, chemotherapy, radiotherapy, cytokine, bone marrow, stem cell transplant.



PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents
 United States Patent and Trademark
 Office
 Box PCT
 Washington, D.C. 20231
 ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 12 May 2000 (12.05.00)	
International application No. PCT/US99/18120	Applicant's or agent's file reference 012712-738
International filing date (day/month/year) 11 August 1999 (11.08.99)	Priority date (day/month/year) 11 August 1998 (11.08.98)
Applicant GRILLO-LÓPEZ, Antonio	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
 13 March 2000 (13.03.00)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Claudio Borton Telephone No.: (41-22) 338.83.38
---	--

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561

562

563

564

565

566

567

568

569

570

571

572

573

574

575

576

577

578

579

580

581

582

583

584

585

586

587

588

589

590

591

592

593

594

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612

613

614

615

616

617

618

619

620

621

622

623

624

625

626

627

628

629

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644

645

646

647

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

676

677

678

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

738

739

740

741

742

743

744

745

746

747

748

749

750

751

752

753

754

755

756

757

758

759

760

761

762

763

764

765

766

767

768

769

770

771

772

773

774

775

776

777

778

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

810

811

812

813

814

815

816

817

818

819

820

821

822

823

824

825

826

827

828

829

830

831

832

833

834

835

836

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

857

858

859

860

861

862

863

864

865

866

867

868

869

870

871

872

873

874

875

876

877

878

879

880

881

882

883

884

885

886

887

888

889

890

891

892

893

894

895

896

897

898

899

900

901

902

903

904

905

906

907

908

909

910

911

912

913

914

915

916

917

918

919

920

921

922

923

924

925

926

927

928

929

930

931

932

933

934

935

936

937

938

939

940

941

942

943

944

945

946

947

948

949

950

951

952

953

954

955

956

957

958

959

960

961

962

963

964

965

966

967

968

969

970

971

972

973

974

975

976

977

978

979

980

981

982

983

984

985

986

987

988

989

990

991

992

993

994

995

996

997

998

999


1000

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)



Applicant's or agent's file reference 012712-738	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US99/18120	International filing date (day/month/year) 11 AUGUST 1999	Priority date (day/month/year) 11 AUGUST 1998
International Patent Classification (IPC) or national classification and IPC IPC(7): A61K 39/395, 38/00; C07K 16/00 and US Cl.: 424/130.1; 514/2; 530/387.1		
Applicant IDEC PHARMACEUTICALS CORPORATION		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
2. This REPORT consists of a total of 4 sheets.
- ☐ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 0 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of report with regard to novelty, inventive step or industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 13 MARCH 2000	Date of completion of this report 29 SEPTEMBER 2000
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer <i>[Signature]</i> SUSAN LUGAR
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0916



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/18120

I. Basis of the report**1. With regard to the elements of the international application:***☒ the international application as originally filed☒ the description:

pages 1-43

pages NONE

pages NONE

, as originally filed

, filed with the demand

, filed with the letter of

☒ the claims:

pages 44-47

pages NONE

pages NONE

pages NONE

, as originally filed

, as amended (together with any statement) under Article 19

, filed with the demand

, filed with the letter of

☒ the drawings:

pages NONE

pages NONE

pages NONE

, as originally filed

, filed with the demand

, filed with the letter of

☒ the sequence listing part of the description:

pages NONE

pages NONE

pages NONE

, as originally filed

, filed with the demand

, filed with the letter of

2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language _____ which is:

☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).☐ the language of publication of the international application (under Rule 48.3(b)).☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).**3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:**☐ contained in the international application in printed form.☐ filed together with the international application in computer readable form.☐ furnished subsequently to this Authority in written form.☐ furnished subsequently to this Authority in computer readable form.☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.**4. ☒ The amendments have resulted in the cancellation of:**☒ the description, pages NONE☒ the claims, Nos. NONE☒ the drawings, sheets/fig. NONE**5. ☒ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).****

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

** Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/18120

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**1. statement**

Novelty (N)	Claims <u>3,6,12-16,18,27-31</u>	YES
	Claims <u>1,2,4,5,7-11,17,19-26</u>	NO
Inventive Step (IS)	Claims <u>12-16</u>	YES
	Claims <u>1-11, 17-31</u>	NO
Industrial Applicability (IA)	Claims <u>1-31</u>	YES
	Claims <u>NONE</u>	NO

2. citations and explanations (Rule 70.7)

Claims 1,2,4,5,7-11,17,19-26 lack novelty under PCT Article 33(2) as being anticipated by US 5,736137.

US 5,736137 teaches treating patients with relapsed B cell lymphoma with a single dose or multiple dose at weekly interval of anti-CD20 antibody C2B8 (columns 28-29). Thus the prior art teaching is the same as the method of claims 1, and 2, because in multiple dose therapy, the patient was treated previously with an anti-CD20 antibody during a second treatment.

US 5,736137 teaches that, as alternative therapy strategies, a radiolabeled C2B8 is used first, followed by single or multiple treatment with C2B8 (column 31). US 5,736137 also teaches the use of chemotherapeutic agent preceding or following treatment with C2B8 (columns 31-32). C2B8 is a chimeric anti-CD20 antibody (column 13, example F). Thus the prior art method is the same as the method of claims 4, 5, and 17-20.

US 5,736137 teaches the use of a therapeutic dose of C2B8 followed within about one week with a combination of C2B8 and radiolabeled anti-CD20 antibody Y2B8 (column 31). C2B8 is a chimeric anti-CD20 antibody (column 13, example F). Thus the method taught by prior art is the same as the methods of claims 7-11, i.e. administration of a chimeric anti-CD20, followed in a week with administration of radiolabeled anti-CD20 antibody Y2B8.

US 5,736137 teaches treatment with radiolabeled anti-CD20 antibody Y2B8, and stem cell or bone marrow transplantation (column 19, clinical analysis of Y2B8 therapy). US 5,736137 also teaches that administration of an anti-CD20 antibody would reduce the level of CD20 positive cells as compared to the level of CD20 positive cells in untreated animals. Thus the method taught by prior art is the same as the methods in claims 21-23. Furthermore, the method taught by prior art is inherently the same as the method of claims 24-26, because administration of an anti-CD20 antibody would inherently reduce the level of CD20 positive cells.

(Continued on Supplemental Sheet.)



INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/18120

Supplemental Box

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of: Boxes I - VIII

Sheet 10

I. BASIS OF REPORT:

5. (Some) amendments are considered to go beyond the disclosure as filed:
NONE

V. 2. REASONED STATEMENTS - CITATIONS AND EXPLANATIONS (Continued):

Claims 1-11, 17-31 lack an inventive step under PCT Article 33(3) as being obvious over US 5,736,137 in view of Muzaffar et al.

The teaching of US 5,736,137 has been set forth. Muzaffar et al teach that 61% of Non-Hodgkin's lymphoma are of B-cell phenotype (p.108), and that all different grades of Non-Hodgkin's lymphoma express CD20, as detected by the marker L26 for CD20 (table). Therefore, it would have been obvious to treat Non-Hodgkin's lymphoma, using the method of US 5,736,137 because of the following reasons: 1) 61% of Non-Hodgkin's lymphoma are of B-cell phenotype (p.108), 2) all different grades of Non-Hodgkin's lymphoma express CD20, and 3) anti-CD20 antibody targeting CD20 is effective in treating B-cell lymphoma.

----- NEW CITATIONS -----

NONE





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 39/395, 38/00, C07K 16/00	A1	(11) International Publication Number: WO 00/09160 (43) International Publication Date: 24 February 2000 (24.02.00)
(21) International Application Number: PCT/US99/18120 (22) International Filing Date: 11 August 1999 (11.08.99) (30) Priority Data: 60/096,180 11 August 1998 (11.08.98) US (71) Applicant (for all designated States except US): IDEC PHARMACEUTICALS CORPORATION [US/US]; 11011 Torreyana Road, San Diego, CA 92121 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): GRILLO-LÓPEZ, Antonio [US/US]; P.O. Box 3797, Rancho Santa Fe, CA 92067 (US). (74) Agents: GESS, E., Joseph et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: COMBINATION THERAPIES FOR B-CELL LYMPHOMAS COMPRISING ADMINISTRATION OF ANTI-CD20 ANTI-BODY		
(57) Abstract New combined therapeutic regimens for treatment of B-cell lymphomas are disclosed which comprise in particular administration of anti-CD20 antibodies to patients having low-, intermediate- or high-grade non-Hodgkins lymphomas.		



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



-1-

**Combination Therapies for B-Cell Lymphomas
Comprising Administration of Anti-CD20 Antibody**

Field of the Invention

The invention relates to the use of anti-CD20 antibodies or fragments thereof in the treatment of B-cell lymphomas, particularly the use of such antibodies and fragments in combined therapeutic regimens.

Background of the Invention

The use of antibodies to the CD20 antigen as diagnostic and/or therapeutic agents for B-cell lymphoma has previously been reported. CD20 is a useful marker or target for B-cell lymphomas as this antigen is expressed at very high densities on the surface of malignant B-cells, i.e., B-cells wherein unabated proliferation can lead to B-cell lymphomas.

CD20 or Bp35 is a B-lymphocyte-restricted differentiation antigen that is expressed during early pre-B-cell development and remains until plasma cell differentiation. It is believed by some that the CD20 molecule may regulate a step in the B-cell activation process which is required for cell cycle initiation and differentiation. Moreover, as noted, CD20 is usually expressed at very high levels on neoplastic ("tumor") B-cells. The CD20 antigen is appealing for targeted therapy, because it does not shed, modulate, or internalize.

Previous reported therapies involving anti-CD20 antibodies have involved the administration of a therapeutic anti-CD20 antibody either alone or in conjunction with a second radiolabeled anti-CD20 antibody, or a chemotherapeutic agent.

In fact, the Food and Drug Administration has approved the therapeutic use of one such anti-CD20 antibody, Rituxan®, for use in relapsed and previously treated low-grade non-Hodgkin's lymphoma (NHL). Also, the use of Rituxan®

-2-

in combination with a radiolabeled murine anti-CD20 antibody has been suggested for the treatment of B-cell lymphoma.

However, while anti-CD20 antibodies and, in particular, Rituxan® (U.S.; in Britain, MabThera®; in general Rituximab®), have been reported to be effective for treatment of B-cell lymphomas, such as non-Hodgkin's lymphoma, the treated patients are often subject to disease relapse. Therefore, it would be beneficial if more effective treatment regimens could be developed.

More specifically, it would be advantageous if anti-CD20 antibodies had a beneficial effect in combination with other lymphoma treatments, and if new combined therapeutic regimens could be developed to lessen the likelihood or frequency of relapse. Also, it would be helpful if current treatment protocols for B-cell lymphoma were improved whereby patients with lymphomas which are refractory to other treatment methods could be treated with chimeric or radiolabeled anti-CD20 antibodies. It would also be helpful if treatment with anti-CD20 antibodies, particularly in combination with other treatments, could be used as therapy for other types of lymphoma besides low grade, follicular non-Hodgkins lymphoma (NHL).

Summary of the Invention

The present invention discloses combined therapeutic treatments for B-cell lymphomas, and reports the benefits of treating relapsed or refractory B-cell lymphomas with chimeric and radiolabeled anti-CD20 antibodies. In particular, it has been found that treatment with anti-CD20 antibody provides a beneficial synergistic effect when administered in combination with cytokines, radiotherapy, myeloablative therapy, or chemotherapy. Surprisingly, patients who had prior bone marrow or stem cell transplantation had an unexpected increase in the overall response rate when compared with patients with no prior therapy.

-3-

Detailed Description of the Invention

This invention encompasses combined therapeutic regimens for the treatment of B-cell lymphomas. In general, such methods include a method for treating relapsed B-cell lymphoma, where a patient having prior treatment for lymphoma has relapsed and is administered a therapeutically effective amount of a chimeric anti-CD20 antibody. Such prior treatments can include, for example, previous treatment with anti-CD20 antibodies, treatments which included a bone marrow or stem cell transplantation, radiotherapy and chemotherapy. The previous chemotherapy may be selected from a wide group of chemotherapeutic agents and combination regimens, including CHOP, ICE, Mitozantrone, Cytarabine, DVP, ATRA, Idarubicin, hoelzer chemotherapy regime, La La chemotherapy regime, ABVD, CEOP, 2-CdA, FLAG & IDA with or without subsequent G-CSF treatment), VAD, M & P, C-Weekly, ABCM, MOPP and DHAP.

Also included in the methods of the invention are methods for treating a subject having B-cell lymphoma wherein the subject is refractory for other therapeutic treatments, including all those listed above, i.e., treatment with chimeric anti-CD20 antibody, treatments which included a bone marrow or stem cell transplantation, radiotherapy and chemotherapy. In particular, encompassed are methods of treating a patient who has not exhibited appreciable tumor remission or regression after administration of a chimeric anti-CD20 antibody, comprising administering to said patient a radiolabeled anti-CD20 antibody.

In particular, the methods of treating a patient with a radiolabeled antibody after a chimeric antibody are performed whereby the radiolabeled anti-CD20 antibody is administered from about one week to about two years after said administration of said chimeric anti-CD20 antibody. More particularly, the

-4-

radiolabeled anti-CD20 antibody is administered from about one week to about nine months after said administration of said chimeric anti-CD20 antibody.

While any anti-CD20 antibodies can be used for the methods of the present invention, a preferred chimeric antibody is C2B8 (IDEC Pharmaceuticals, Rituximab®). A preferred radiolabeled antibody is Y2B8, which is a murine antibody labeled with yttrium-90 (⁹⁰Y). However, antibodies with other radiolabels may be used, particularly those labeled with a beta or alpha isotope. Anti-CD19 antibodies may also be used.

One of skill in the art would know the parameters for choosing a particular type of anti-CD20 antibody. For instance, chimeric and humanized antibodies are beneficial for decreased immunogenicity, and for facilitating antibody effector mediated immune reactions via the human constant region domains. Murine and other mammalian antibodies, in contrast, are beneficial for delivering a radiolabel to the tumor cell, as such antibodies generally have a decreased half-life *in vivo*.

Antibody treatments performed initially to which patients are refractory or have relapsed may include initial treatments with chimeric antibodies or mammalian antibodies. Also encompassed are initial treatments with other antibodies, including anti-CD19 antibodies and anti-Lym antibodies, and treatments with antibodies labeled with cytotoxic moieties, such as toxins, and radiolabels, e.g., Oncolym® (Techniclone) or Bexxar (Coulter).

It should be clear that the combined therapeutic regimens of the present invention can be performed whereby said therapies are given simultaneously, i.e., the anti-CD20 antibody is administered concurrently or within the same time frame (i.e., the therapies are going on concurrently, but the agents are not administered precisely at the same time). The anti-CD20 antibodies of the present invention may also be administered prior to or subsequent to the other therapies.

Sequential administration may be performed regardless of whether the patient responds to the first therapy to decrease the possibility of remission or relapse.

The combined therapies of the present invention include a method for treating B-cell lymphoma comprising administering at least one chimeric anti-
5 CD20 antibody and at least one cytokine. In particular, the invention includes a method for treating B-cell lymphoma comprising administering a synergistic therapeutic combination comprising at least one anti-CD20 antibody and at least one cytokine, wherein the therapeutic effect is better than the additive effects of either therapy administered alone. Preferred cytokines are selected from the group
10 consisting of alpha interferon, gamma interferon, IL-2, GM-CSF and G-CSF. Again, the anti-CD20 antibody and the cytokine(s) may be administered sequentially, in either order, or in combination.

Also included in the present invention is a method for treating B-cell lymphoma comprising administering to a patient a therapeutically effective
15 amount of a chimeric anti-CD20 antibody before, during or subsequent to a chemotherapeutic regimen. Such a chemotherapy regimen may be selected from the group consisting of, at the very least, CHOP, ICE, Mitozantrone, Cytarabine, DVP, ATRA, Idarubicin, hoelzer chemotherapy regime, La La chemotherapy regime, ABVD, CEOP, 2-CdA, FLAG & IDA with or without subsequent G-CSF
20 treatment), VAD, M & P, C-Weekly, ABCM, MOPP and DHAP.

Also encompassed are methods for treating B-cell lymphoma comprising administering to a patient a therapeutically effective amount of a chimeric anti-CD20 antibody before, during or subsequent to a bone marrow or peripheral stem cell transplant. Such bone marrow transplant may also be accompanied by other
25 therapeutic regimens such as chemotherapy. The antibodies of the present invention may also be used in a method of reducing residual CD20+ tumor cells in bone marrow or stem cells before or after myeloablative therapy by

-6-

administering to a patient a chimeric anti-CD20 antibody. It may also be possible to use such antibodies *in vitro* to induce apoptosis of tumor cells and reduce or cure bone marrow or stem cell preparations of residual tumor cells before they are infused back into the patient.

5 It should be understood that stem cell transplants may be allogeneic or autologous. If the transplant is allogeneic, i.e., from another person, the disclosed therapeutic regimens may include treatments with immunosuppressive drugs before administration of the anti-CD20 antibodies. Co-administration of other
10 drugs designed to enhance acceptance of the transplant and stimulate the production and differentiation of immune cells is also contemplated. For instance, it has been shown that administration of GM-CSF to marrow transplant recipients promotes the development of specific bone marrow cells which in turn produces circulating infection-fighting neutrophils, and increased the survival rate of marrow transplant recipients.

15 The methods of the present invention may be used to treat a variety of B-cell lymphomas, including low grade/ follicular non-Hodgkin's lymphoma (NHL), small lymphocytic (SL) NHL, intermediate grade/ follicular NHL, intermediate grade diffuse NHL, high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL and
20 Waldenstrom's Macroglobulinemia. It should be clear to those of skill in the art that these lymphomas will often have different names due to changing systems of classification, and that patients having lymphomas classified under different names may also benefit from the combined therapeutic regimens of the present invention.

25 For instance, a recent classification system proposed by European and American pathologists is called the Revised European American Lymphoma (REAL) Classification. This classification system recognizes Mantle cell

-7-

lymphoma and Marginal cell lymphoma among other peripheral B-cell neoplasms, and separates some classifications into grades based on cytology, i.e., small cell, mixed small and large, large cell. It will be understood that all such classified lymphomas may benefit from the combined therapies of the present invention.

5 The U.S. National Cancer Institute (NCI) has in turn divided some of the REAL classes into more clinically useful "indolent" or "aggressive" lymphoma designations. Indolent lymphomas include follicular cell lymphomas, separated into cytology "grades," diffuse small lymphocytic lymphoma/chronic lymphocytic leukemia (CLL), lymphoplasmacytoid/Waldenstrom's Macroglobulinemia,
10 Marginal zone lymphoma and Hairy cell leukemia. Aggressive lymphomas include diffuse mixed and large cell lymphoma, Burkitt's lymphoma/diffuse small non-cleaved cell lymphoma, Lymphoblastic lymphoma, Mantle cell lymphoma and AIDS-related lymphoma. These lymphomas may also benefit from the combined therapeutic regimens of the present invention.

15 Non-Hodgkin's lymphoma has also been classified on the basis of "grade" based on other disease characteristics including low-grade, intermediate-grade and high-grade lymphomas. Low-grade lymphoma usually presents as a nodal disease, and is often indolent or slow-growing. Intermediate- and high-grade disease usually presents as a much more aggressive disease with large extranodal
20 bulky tumors. Intermediate- and high-grade disease, as well as low grade NHL, may benefit from the combined therapeutic regimens of the present invention.

 The Ann Arbor classification system is also commonly used for patients with NHL. In this system, stages I, II, III, and IV of adult NHL can be classified into A and B categories depending on whether the patient has well-defined
25 generalized symptoms (B) or not (A). The B designation is given to patients with the following symptoms: unexplained loss of more than 10% body weight in the

-8-

6 months prior to diagnosis, unexplained fever with temperatures above 38°C and drenching night sweats. Occasionally, specialized staging systems are used:

Stage I - involvement of a single lymph node region or localized involvement of a single extralymphatic organ or site.

- 5 Stage II - involvement of two or more lymph node regions on the same side of the diaphragm or localized involvement of a single associated extralymphatic organ or site and its regional lymph nodes with or without other lymph node regions on the same side of the diaphragm.

- 10 Stage III - involvement of lymph node regions on both sides of the diaphragm, possibly accompanying localized involvement of an extralymphatic organ or site, involvement of the spleen, or both.

Stage IV - disseminated (multi-focal) involvement of 1 or more extralymphatic sites with or without associated lymph node involvement or isolated extralymphatic organ involvement with distant (non-regional) nodal involvement.

- 15 For further details, see The International Non-Hodgkin's Lymphoma Prognostic Factors Project: A predictive model for aggressive non-Hodgkin's lymphoma. *New England J. Med.* 329(14): 987-994 (1993).

Preferred antibodies, dosage regimens and particular combinations of therapy will now be illustrated by way of the following exemplary data.

20 **Rituximab® and Y2B8**

Non-Hodgkin's lymphoma (NHL) affects approximately 250,000 people in the United States. The majority of patients with NHL are not cured by chemotherapy, radiotherapy, or high-dose treatment with autologous bone marrow (ABMT) or peripheral blood stem cell (PBSC) support.

- 25 Approximately 80% of non-Hodgkin's lymphomas are B-cell malignancies and > 95% of these express the CD20 antigen on the cell surface. This antigen is an attractive target for immunotherapy because it is found exclusively on B-cells,

-9-

and not on hematopoietic stem cells, pro-B cells, normal plasma cells, or other normal tissues. It is not shed from the cell surface and does not modulate upon antibody binding (1).

Rituximab® is one of a new generation of monoclonal antibodies
5 developed to overcome limitations encountered with murine antibodies, including short half-life, limited ability to stimulate human effector functions, and immunogenicity (2,3).

Rituximab® is a genetically engineered monoclonal antibody with murine light-and heavy-chain variable regions and human gamma I heavy-chain and
10 kappa light-chain constant regions. The chimeric antibody is composed of two heavy chains of 451 amino acids and two light chains of 213 amino acids and has an approximate molecular weight of 145 kD. Rituximab® is more effective than its murine parent in fixing complement and mediating ADCC, and it mediates CDC in the presence of human complement (4). The antibody inhibits cell growth
15 in the B-cell lines FL-18, Ramos, and Raji, sensitizes chemoresistant human lymphoma cell lines to diphtheria toxin, ricin, CDDP, doxorubicin, and etoposide, and induces apoptosis in the DHL-4 human B-cell lymphoma line in a dose-dependent manner (5). In humans, the half-life of the antibody is approximately 60 hours after the first infusion and increases with each dose to 174 hours after the
20 fourth infusion. The immunogenicity of the antibody is low; of 355 patients in seven clinical studies, only three (<1%) had a detectable anti-chimeric antibody (HACA) response.

Rituximab® was genetically engineered using the murine 2B8 antibody. The 2B8 antibody has also been conjugated to different radiolabels for diagnostic
25 and therapeutic purposes. To this end, copending application Serial Nos. 08/475,813, 08/475,815 and 08/478,967, herein incorporated by reference in their entirety, disclose radiolabeled anti-CD20 conjugates for diagnostic "imaging" of

-10-

B-cell lymphoma tumors before administration of therapeutic antibody. "In2B8" conjugate comprises a murine monoclonal antibody, 2B8, specific to human CD20 antigen, that is attached to Indium[111] (^{111}In) via a bifunctional chelator, i.e., MX-DTPA (diethylenetriaminepentaacetic acid), which comprises a 1:1 mixture
5 of 1-isothiocyanatobenzyl-3-methyl-DTPA and 1-methyl-3-isothiocyanatobenzyl-DTPA. Indium-[111] is selected as a diagnostic radionuclide because it emits gamma radiation and finds prior usage as an imaging agent.

Patents relating to chelators and chelator conjugates are known in the art. For instance, U.S. Patent No. 4,831,175 of Gansow is directed to polysubstituted
10 diethylenetriaminepentaacetic acid chelates and protein conjugates containing the same, and methods for their preparation. U.S. Patent Nos. 5,099,069, 5,246,692, 5,286,850, and 5,124,471 of Gansow also relate to polysubstituted DTPA chelates. These patents are incorporated herein in their entirety.

The specific bifunctional chelator used to facilitate chelation in MX-DTPA
15 was selected as it possesses high affinity for trivalent metals, and provides for increased tumor-to-non-tumor ratios, decreased bone uptake, and greater *in vivo* retention of radionuclide at target sites, i.e., B-cell lymphoma tumor sites. However, other bifunctional chelators are known in the art and may also be beneficial in tumor therapy.

20 Also disclosed in U.S. Patent No. 5,736,137 are radiolabeled therapeutic antibodies for the targeting and destruction of B-cell lymphomas and tumor cells. In particular, the Y2B8 conjugate comprises the same anti-human CD20 murine monoclonal antibody, 2B8, attached to yttrium-[90] (^{90}Y) via the same bifunctional chelator. This radionuclide was selected for therapy for several
25 reasons. The 64 hour half-life of ^{90}Y is long enough to allow antibody accumulation by the tumor and, unlike e.g. ^{131}I , it is a pure beta emitter of high energy with no accompanying gamma irradiation in its decay, with a range of 100

-11-

to 1000 cell diameters. The minimal amount of penetrating radiation allows for outpatient administration of ^{90}Y -labeled antibodies. Furthermore, internalization of labeled antibodies is not required for cell killing, and the local emission of ionizing radiation should be lethal for adjacent tumor cells lacking the target antigen.

Because the ^{90}Y radionuclide was attached to the 2B8 antibody using the same bifunctional chelator molecule MX-DTPA, the Y2B8 conjugate possesses the same advantages discussed above, e.g., increased retention of radionuclide at a target site (tumor). However, unlike ^{111}In , it cannot be used for imaging purposes due to the lack of gamma radiation associated therewith. Thus, a diagnostic "imaging" radionuclide, such as ^{111}In , can be used for determining the location and relative size of a tumor prior to and/or following administration of therapeutic chimeric or ^{90}Y -labeled antibodies in the combined regimens of the invention. Additionally, indium-labeled antibody enables dosimetric assessment to be made.

Depending on the intended use of the antibody, i.e., as a diagnostic or therapeutic reagent, other radiolabels are known in the art and have been used for similar purposes. For instance, radionuclides which have been used in clinical diagnosis include ^{131}I , ^{125}I , ^{123}I , ^{99}Tc , ^{67}Ga , as well as ^{111}In . Antibodies have also been labeled with a variety of radionuclides for potential use in targeted immunotherapy (Peirersz et al. (1987) The use of monoclonal antibody conjugates for the diagnosis and treatment of cancer. *Immunol. Cell Biol.* 65: 111-125). These radionuclides include ^{188}Re and ^{186}Re as well as ^{90}Y , and to a lesser extent ^{199}Au and ^{67}Cu . I-(131) has also been used for therapeutic purposes. U.S. Patent No. 5,460,785 provides a listing of such radioisotopes and is herein incorporated by reference.

-12-

As reported in U.S. Patent No. 5,736,137, administration of the radiolabeled Y2B8 conjugate, as well as unlabeled chimeric anti-CD20 antibody, resulted in significant tumor reduction in mice harboring a B-cell lymphoblastic tumor. Moreover, human clinical trials reported therein showed significant B-cell
5 depletion in lymphoma patients infused with chimeric anti-CD20 antibody. In fact, chimeric 2B8 has recently been heralded the nation's first FDA-approved anti-cancer monoclonal antibody under the name of Rituxan®. Thus, at least one chimeric anti-CD20 antibody has been shown to demonstrate therapeutic efficacy in the treatment of B-cell lymphoma.

10 In addition, U.S. Patent No. 5,736,137, herein incorporated by reference, discloses sequential administration of Rituxan®, a chimeric anti-CD20, with both or either indium-labeled or yttrium-labeled murine monoclonal antibody. Although the radiolabeled antibodies used in these combined therapies are murine antibodies, initial treatment with chimeric anti-CD20 sufficiently depletes the B-
15 cell population such that the HAMA response is decreased, thereby facilitating a combined therapeutic and diagnostic regimen.

Thus, in this context of combined immunotherapy, murine antibodies may find particular utility as diagnostic reagents. Moreover, it was shown in U.S. Patent No. 5,736,137 that a therapeutically effective dosage of the yttrium-labeled
20 anti-CD20 antibody following administration of Rituxan® is sufficient to (a) clear any remaining peripheral blood B-cells not cleared by the chimeric anti-CD20 antibody; (b) begin B-cell depletion from lymph nodes; or (c) begin B-cell depletion from other tissues.

Thus, conjugation of radiolabels to cancer therapeutic antibodies provides
25 a valuable clinical tool which may be used to assess the potential therapeutic efficacy of such antibodies, create diagnostic reagents to monitor the progress of treatment, and devise additional therapeutic reagents which may be used to

-13-

enhance the initial tumor-killing potential of the chimeric antibody. Given the proven efficacy of an anti-CD20 antibody in the treatment of non-Hodgkin's lymphoma, and the known sensitivity of lymphocytes to radioactivity, it would be highly advantageous for such chimeric and radiolabeled therapeutic antibodies to find use in combined therapeutic regimens which decrease the frequency of relapsed or refractory non-Hodgkin's lymphoma. In addition, it would be beneficial if such combined therapeutic regimens found use in the treatment of other B-cell lymphomas.

LOW-GRADE OR FOLLICULAR NHL

Single-Agent Studies with Relapsed or Refractory NHL

FDA approval of Rituximab® was based on five single-agent studies primarily in patients with low-grade or follicular NHL. An early Phase I study of single Rituximab® infusions ranging from 10 - 500 mg/m² demonstrated that the maximum tolerated dose had not been reached; however, the length of infusion time at the highest dose was not considered feasible for outpatient therapy. The ORR in 15 patients was 13% (Table 1)(6).

Table 1
Rituximab®: Summary of Efficacy Results

Study Description	Indication	N*	ORR	CR	PR	Median DR (months)	Median TIP (months)	References
Phase I/II, Single-Dose Single Agent	Relapsed B-Cell Lymphoma	15	2 (13%)	0 (0%)	2 (13%)	NA†	8.1	6
Phase I/II, Multiple-Dose Dose-Ranging	Relapsed Low-, Intermediate-, and High-Grade Lymphoma	34	17 (50%)	3 (9%)	14 (41%)	8.6	10.2	7
Phase II; Multiple-Dose Combined with CHOP	Newly Diagnosed and Relapsed Low-Grade or Follicular B-Cell Lymphoma	38	38 (100%)	22 (58%)	16 (42%)	35.3+	36.7+	21,22
Phase III, Multiple-Dose Single-Agent	Relapsed Low-Grade or Follicular B-Cell Lymphoma	151	76 (50%)	9 (6%)	67 (44%)	11.6	13.2	8, 9
Phase II, Multiple-Dose Single-Agent	Relapsed Low-Grade or Follicular B-Cell Lymphoma	35	21 (60%)	5 (14%)	16 (46%)	13.4+	19.4+	13
Phase II, Multiple-Dose, Combined with Interferon	Relapsed Low-Grade or Follicular B-Cell Lymphoma	38	17 (45%)	4 (11%)	13 (34%)	22.3+	25.2+	29
Phase II, Multiple-Dose, Single-Agent	Relapsed Low-Grade or Follicular B-Cell Lymphoma, Bulky Disease	28	12 (43%)	1 (4%)	11 (39%)	5.9	8.1	14

Study Description	Indication	N*	ORR	CR	PR	Median DR (months)	Median TIP (months)	References
Phase II, Multiple-Dose, Single-Agent	Relapsed Low-Grade or Follicular B-Cell Lymphoma, Retreatment	57	23 (40%)	6 (11%)	17 (29%)	15.0+	16.7+	19,20
Phase II, Multiple-Dose Combined with CHOP Modality	Previously Untreated Intermediate- or High-Grade Lymphoma	30	29 (96%)	19 (63%)	10 (33%)	11+	17+	34
Phase II, Alternative Multiple Dosing	Intermediate- or High-Grade B-Cell Lymphoma	54	17 (32%)	5 (9%)	12 (22%)	NA†	8.2+	33

*N = number of evaluable patients

†Not available

-16-

In Phase I of a Phase I/II dose-ranging study, patients received 125 - 375 mg/m² administered as four weekly infusions. No dose-related toxicities were demonstrated, and 375 mg/m² was chosen as the Phase II dose. Tumor regressions were observed in 17 of 37 (46%) patients who received this dose, including 3
5 (8%) complete responses (CR) and 14 (38%) partial responses PR (7).

A subsequent single-arm pivotal study of Rituximab® infused at 375 mg/m² weekly times four was conducted in 166 patients with relapsed or refractory, low-grade or follicular NHL (International Working Formulation [IWF] Types A - D and REAL classification, small lymphocytic lymphoma,
10 Follicular center, follicular Grades I, II, III(8)). Patients with tumor masses > 10 cm or with > 5000 lymphocytes/μL in the peripheral blood were excluded from this study. The median age was 58 years (105 men and 61 women) and the median number of prior treatments was three. Bone marrow involvement was present in 56% of 149 patients evaluated. Forty-five percent had ≥ 2 extranodal
15 sites and 41% had bulky disease (≥ 5 cm).

Complete response required the regression of all lymph nodes to < 1 × 1 cm² demonstrated on two occasions at least 28 days apart on neck, chest, abdomen, and pelvic CT scans, resolution of all symptoms and signs of lymphoma, and normalization of bone marrow, liver, and spleen. Partial response
20 required a ≥ 50% decrease in the sum of the products of perpendicular measurements of lesions without any evidence of progressive disease for at least 28 days. Patients who did not achieve a CR or PR were considered non-responders, even if a net decrease (> 50%) of measurable disease was observed. Time to progression was measured from the first infusion until progression.

25 The overall response rate (ORR) was 48% with a 6% CR and a 42% PR rate(8). The median time to progression (TTP) for responders was 13.2 months

-17-

and the median duration of response (DR) was 11.6 months. Twenty-two of 80 (28%) responders remain in ongoing remission at 20.9+ to 32.9+ months (9).

Administration of Rituximab® resulted in a rapid and sustained depletion of B-cells. Circulating B-cells were depleted within the first three doses with
5 sustained depletion for up to six to nine months post-treatment in 83% of patients. Median B-cell levels returned to normal by 12 months following treatment. Although median NK cell counts remained unchanged, a positive correlation was observed between higher absolute NK cell counts at baseline and response to Rituximab® (10).

10 Several baseline prognostic factors were analyzed to determine their correlation to response. Significantly, in 23 patients relapsed after ABMT or PBSC, the ORR was 78% versus 43% in patients who did not undergo prior high-dose therapy ($p < 0.01$). In a multivariate analysis, the ORR was higher in patients with follicular NHL as compared with small lymphocytic lymphoma (58% vs.
15 12%, $p < 0.01$), and higher in patients with chemosensitive relapse as compared with chemoresistant relapse (53% vs. 36%, $p = 0.06$). No effect on response rate was associated with: age > 60 years, extranodal disease, prior anthracycline therapy, or bone marrow involvement.

A statistically significant correlation was found between the median serum
20 antibody concentration and response at multiple time points during treatment and follow up (11).

Serum levels of antibody were higher in patients with follicular NHL compared with small lymphocytic lymphoma. Mean serum antibody was also inversely correlated with measurements of tumor bulk and with the number of
25 circulating B-cells at baseline. The association of lower serum antibody concentrations with higher numbers of circulating NHL cells and with higher tumor bulk suggest that the main mode of antibody clearance is to tumor cells.

-18-

The association of high serum antibody concentrations with response and lower tumor bulk or circulating cells suggests that higher or more doses of Rituximab® may be necessary to induce responses in some subsets of patients, such as those with bulky disease.

5 Nevertheless, responses were seen with Rituximab® in 43% of patients with tumors > 5 cm and in 35% of patients with tumors > 7 cm, suggesting that treatment of patients with bulky disease with Rituximab® is feasible. This is surprising considering it was long thought that antibody therapy is not conducive to treating bulky disease due to the compact nature of the tumors.

10 In a study conducted in Japan (12), patients with relapsed B-cell lymphoma were treated with either 250 mg/m² (N = 4) or 375 mg/m² (N = 8) of Rituximab® weekly times four. Of 11 evaluable patients, 8 had follicular NHL, 2 had diffuse large-cell NHL, and one had mantle-cell lymphoma. Two of the 11 had a CR and 5 had a PR for an ORR of 64%; all responders had follicular histology.

15 Because Rituximab® serum levels and response were positively correlated in previous studies, a Phase II study of eight weekly doses of 375 mg/m² Rituximab® was conducted in low-grade or follicular NHL patients. The ORR was 60% in evaluable patients, with a 14% CR and a 46% PR rate. Median values for TTP in responders and DR were 13.4+ months and 19.4+ months, respectively
20 (13). Though it is difficult to compare across studies, it appears that TTP and DR may be improved by using more doses.

 Contrary to early assumptions about antibody therapy being useful only in micrometastatic disease, Rituximab® is quite active in high bulk disease. In a separate study, 31 patients with relapsed or refractory, bulky low-grade NHL
25 (single lesion of > 10 cm in diameter) received 375 mg/m² Rituximab® as four weekly infusions. Twelve of 28 evaluable patients (43%) demonstrated a CR (1, 4%) or PR (11, 39%)(14).

Waldenstrom's Macroglobulinemia

Waldenstrom's Macroglobulinemia (WM) is a malignancy wherein B lymphocytes secrete excessive amounts of IgM antibodies. WM usually occurs in people over sixty, but has been detected in adults in their early thirties. WM today is considered a rare incurable indolent malignancy, which has in the past been treated by plasmaphoresis to reduce serum viscosity. Chemotherapeutic drugs such as an alkylating agent and a corticosteroid are often prescribed. The most recommended drug for WM has been Leustatin (2CdA).

A report on seven patients with Waldenstrom's macroglobulinemia where the patients were treated with Rituximab® (375 mg/m² weekly times 4)(15) noted responses in 4 (57%) of patients. Median progression-free survival was 8 months (range 3 - 27+ months). Thus, Rituximab® should be useful in combined therapeutic protocols, particularly with chemotherapeutic reagents such as 2CdA.

Chronic Lymphocytic Leukemia (CLL)

CLL is the liquid (leukemic) equivalent of small lymphocytic lymphoma (SLL). Patients with SLL had lower serum levels and a lower response rate when treated with the standard dose of Rituximab® than patients with other low-grade NHL subtypes. This is probably due to the very high levels of circulating tumor cells in patients with CLL, and because malignant cells involved in CLL are thought to have reduced levels of expression of CD20 on the cell surface.

Nevertheless, the present inventors have discovered that hematologic malignancies such as CLL may be treated with Rituximab®. A recent clinical study evaluated treatment of CLL patients at higher doses of Rituximab® (16). All patients receive a first dose of 375 mg/m³ to minimize infusion-related side effects. Subsequent weekly dosages (3) remained the same but were given at an increased dose level. Sixteen patients have been treated at dosages of 500-1500 mg/m³. Median age was 66 years (range, 25-78). Eighty-one percent had end-

-20-

stage III-IV disease. Medium white blood cell count was $40 \times 10^9/L$ (range, 4-200), Hgb 11.6 g/dl (range, 7.7-14.7), platelets $75 \times 10^9/L$ (range, 16-160), median β_2 immunoglobulin was 4.5 mg/L (range, 3.1-9.2). Median numbers of prior therapies was 2.5 (range 1-9). Sixty percent of patients were refractory to treatment. Two patients developed severe hypertension with the first dose (375 mg/m³); another one received further therapy. Toxicity at subsequent escalated dosages has been mild although no patient at the 1500 mg/m³ dose level has been fully evaluated. Eight patients have completed therapy (4 at 500 mg/m³, 3 at 650 mg/m³, 1 at 825 mg/m³). One patient treated at 560 mg/m³ achieved full remission. One patient has progressive lymphocytosis on treatment and all other patients had reduction in peripheral blood lymphocytosis but less effect on lymph nodes. Dose escalation studies are ongoing.

Another approach to improving response in CLL patients is to upregulate the CD20 antigen using cytokines. In an *in vitro* study, mononuclear cells from CLL patients were incubated for 24 hours with various cytokines. Flow cytometry results showed significant up-regulation by IL-4, GM-CSF, and TNF-alpha (17). In fact, recent data suggests that the upregulation of CD20 observed on CLL cells may be limited to tumor cells (Venogopal et al. Poster - PanPacific Lymphoma meeting, June 1999. Cytokine-induced upregulation of CD20 antigen expression in chronic lymphocytic leukemia (CLL) cells may be limited to tumor cells). Preliminary data also suggest that interferon alpha also upregulates CD20 on CLL cells after only 24 hours when applied at a concentration of 500 to 1000 U/ml.

Thus, by administering certain cytokines to CLL patients prior to or concurrently with administration of Rituximab®, the expression of CD20 on the surface of malignant B-cells may be upregulated, thereby rendering CD20, as well as other cell surface markers such as CD19, a more attractive target for immunotherapy. A collaborative study has been initiated to test for optimal

-21-

cytokine doses for CD20 upregulation in vivo. The study protocol involves treating ten patients initially with GM-CSF at 250 mcg/m² SQ QD X 3, ten patients with IL-4 mcg/kg SQ QD X 3, and ten patients with G-CSF at 5 mcg/kg SQ QD X 3. Mononuclear cells will be separated by Ficon Hypaque centrifugation for apoptotic studies to determine if upregulation of CD20 translates to enhanced killing of tumor cells by Rituximab®.

Antibody treatment of CLL can be combined with other conventional chemotherapeutic treatments known to be useful for the treatment of CLL. The most frequently used single agent for CLL is chlorambucil (leukeran), given either as 0.1 mg/kg daily or 0.4 to 1.0 mg/kg every 4 weeks. Chlorambucil is often combined with oral prednisone (30 to 100 mg/m²/d), which is useful in the management of autoimmune cytopenias. Cyclophosphamide is an alternative to chlorambucil, the usual dose being 1-2 g/m² every 3-4 weeks together with vincristine and steroids (e.g., COP regimen).

Various drug combinations have been used for CLL, including COP (cyclophosphamide, Oncovin, and prednisone), and CHOP (these three drugs plus doxorubicin). Fludarabine has shown an effect in the treatment of CLL, and gave an ORR of 50% in a group of patients treated with 25-30 mg/m²/d every 3-4 weeks. Although some patients have been shown to be refractory for fludarabine. Such patients may also be resistant to 2-CdA because often, patients who are refractory to fludarabine are also refractory to 2-CDA (O'Brien et al. N. Engl. J. Med. 330: 319-322 (1994)).

Hence, anti-CD20 antibody therapy will be particularly useful for patients who are refractory or who have relapsed after treatment with chemotherapeutic drugs. Rituximab® therapy may also be combined with radiotherapy in these patients. TBI with a low fraction size of 15 cGy to total doses of 75 to 150 cGy has been shown to be effective in about one-third of patients.

-22-

A Phase II trial is currently being conducted by CALGB in CLL patients. Rituximab® and fludarabine are administered concurrently, followed by Rituximab® consolidation versus fludarabine induction followed by Rituximab®.

Rituximab® with Myeloablative Therapy

5 Myeloablative therapy has yielded responses in indolent lymphomas; however, residual tumor cells may remain despite high-dose therapy and the PBSC reinfused may contain tumor cells. Rituximab® is being used before stem cell mobilization and after transplant to reduce residual CD20+ tumor cells and contamination of the bone marrow or stem cells harvested. Interim results
10 demonstrated that no CD20+ cells were detectable in harvested cells. Eighteen of 24 patients achieved engraftment and the treatment was well tolerated. PCR testing is ongoing to evaluate residual tumor cells (18).

Retreatment of Relapsed Low-Grade NHL with Rituximab®

A trial evaluating retreatment of 53 patients who had responded to
15 Rituximab® and later relapsed has been reported (19). Seven of 56 evaluable patients (13%) obtained a CR and 16 a PR (29%), for an ORR of 42%. Four patients who had a second response received a third treatment; 3 of these responded.

After treatment with two courses of Rituximab®, one patient's tumor,
20 initially classified as follicular, small cleaved cell NHL, no longer expressed the CD20 antigen and was unresponsive to Rituximab® at the time of transformation to diffuse, large-cell NHL (20).

Thus, while retreatment with Rituximab® is effective for treating patients who have relapsed after prior treatment with Rituximab®, there may be an
25 increased incidence of CD20- tumor cells after secondary treatment. This observation supports the utility of the combined therapeutic treatment regimens described herein.

Combination of Rituximab® and CHOP Chemotherapy for Low-Grade NHL

Chemotherapy with cyclophosphamide, doxorubicin, vincristine, and prednisone (CHOP) is an effective first-line therapy for low-grade or follicular NHL. Though initial response rates are high, relapse eventually occurs and
5 subsequent chemotherapy regimens produce remissions with shorter durations. A Phase II trial was initiated to evaluate the combination of CHOP and Rituximab® (21) in newly diagnosed and relapsed low-grade or follicular NHL because their mechanisms of action are not cross-resistant, and Rituximab® is synergistic with certain cytotoxic drugs, including doxorubicin (5).

10 Twenty-nine of 38 patients received no prior anticancer therapy. CHOP was administered at standard doses every three weeks for six cycles with six infusions of Rituximab® (375 mg/m²). Rituximab® infusions 1 and 2 were administered on Days 1 and 6 before the first CHOP cycle, which started on Day 8. Rituximab® infusions 3 and 4 were given 2 days before the third and fifth
15 CHOP cycles, respectively, and infusions 5 and 6 were given on Days 134 and 141, respectively, after the sixth CHOP cycle.

In this combination study, 100% of the 38 patients treated responded (CR, 58%; PR, 42%). Of 35 evaluable patients who completed treatment, there were 63% CR, and 37% PR(21). Median DR is 35.3+ months with median progression-
20 free survival not reached after a median observation time of 36.7+ months. Twenty patients are still in remission after 36+ months to 53.4+ months (22). This DR is impressive even for first-line treatment, and 24% of this trial population had relapsed after chemotherapy.

In a study to be conducted by CALGB, 40 patients with low-grade NHL
25 will receive Rituximab® weekly times 8 and oral cyclophosphamide daily starting on Day 8. Twenty patients will receive Rituximab® alone for 8 weekly doses.

-24-

A Phase III study conducted by ECOG in patients with low-grade NHL is comparing the combination of cyclophosphamide and fludarabine (Arm A) with standard CVP therapy (Arm B). In the randomization to Arm A or Arm B, patients are stratified by age, tumor burden, histology, and B symptoms. Responders in both arms will undergo a second randomization to Rituximab® maintenance therapy (375 mg/m² weekly times 4 every 6 months for 2 years (Arm C) or to observation (Arm D).

Combination of Rituximab® with Cytokines

Rituximab® plus interferon alpha

Interferon is a cytokine involved in modulating the immune system (23). Mechanisms by which interferon may increase the effectiveness of antibodies include the potentiation of antigen expression (24), increased targeting of antibodies into tumors (25,26), and enhanced cytotoxicity of immunotoxins (27).

In a combination trial, interferon-alpha (Roferon-A), a cytokine with a single-agent clinical activity in NHL (28), and Rituximab® were given to patients with relapsed low-grade or follicular NHL. Interferon-alpha (2.5 or 5 MIU) was administered subcutaneously, three times weekly for 12 weeks. Rituximab® was administered by IV infusion weekly for four doses (375 mg/m²) starting on the fifth week of treatment. The ORR was 45% (17/38 patients); 11% had a CR and 34% had a PR. Kaplan-Meier estimates of the median DR and TTP in responders were 22.3+ and 25.2+ months, respectively (29). Previous combination studies of interferon-alpha and chemotherapeutic regimens containing anthracyclines yielded prolonged time to progression, but did not consistently increase response or survival rates (30-32). These early results suggest that the combination of Rituximab® and interferon-alpha may prolong the time to progression relative to Rituximab® alone.

-25-

Rituximab® plus G-CSF

In a separate study, Rituximab® and G-CSF are being evaluated in relapsed low-grade NHL. It has been demonstrated in vitro as well as in vivo in healthy volunteers that G-CSF, via its effect on myeloid precursor cells, induces FcRI-positive neutrophils that are capable of functioning as effector cells in ADCC. Therefor, a Phase I/II study was initiated to evaluate the toxicity and efficacy of the combined treatment.

Both in Phase I and Phase II, patients were administered a standard dose of G-CSF (5 µg/kg/day) administered for three days, starting 2 days before administration of Rituximab®. Phase I consisted of a dose escalation of Rituximab® (125, 250, or 375 mg/m² weekly X4). Early results in 9 patients evaluated so far yielded an ORR of 67% (44% CR, 22% PR) with minor toxicity in 8 of the 9 patients (33). The most frequent adverse events were fever (4/8 patients), rhinitis (4/8), chills (3/8) and headaches (3/8), which were comparable to the adverse events observed previously in administration of Rituximab® alone. The Phase II part of the study has been initiated, which will examine the efficacy of the combination of G-CSF and 375 mg/m² Rituximab® X4.

Rituximab® plus IL-2

High-dose therapy with autologous peripheral blood stem cells (PBSC) or bone marrow (BM) rescue has been used to treat NHL, however success remains limited by the high risk of relapse, which is 50-80%. In an effort to improve durable remissions post-transplant, immunotherapy including high dose and low dose therapy with IL-2 has been studied in a number of treatment centers. Such studies have suggested that IL-2 therapy does demonstrate early post-transplant anti-Tumor activity.

Initially following autologous transplant, patients display delayed immune reconstitution which potentially results in diminished immune-mediated tumor

-26-

eradication (43, 44). Indeed, it has been shown that both CD4⁺ T cells and cytotoxic CD8⁺ T cells are depressed (45-49). In vitro assays have demonstrated a profound suppression of T cell cytolytic and proliferative responses as well as decreased production of IL-2 in response to mitogens and soluble antigens.

5 However, soluble IL-2 is able to restore these immune responses suggesting that immune cells in patients after autologous transplant are capable of responding to exogenous IL-2 (47). Peripheral blood NK activity also remains lower following BMT than control values and the NK activity is also augmented by addition of exogenous IL-2 (49). These data suggest that administration of IL-2 to patients

10 shortly after stem cell transplant may enhance immune responsiveness at a critical period when tumor burden is minimal and when immune responsiveness in the absence of IL-2 is lacking.

For instance, Caligiuri et al. have shown that IL-2 (Hoffman-LaRoche) administered at 0.45×10^6 U/M²/day by 24 hour CIV for 12 weeks was able to

15 expand the absolute number of CD56 bright NK cells (50-52). This regimen was administered to non-transplant patients in the outpatient setting with little toxicity.

Animal models have shown that non-LAK inducing low doses of IL-2 dramatically enhances anti-tumor activity when administered with tumor-specific T effector cells (53). In addition, Soiffer et al. (54) administered low doses of IL-

20 2 to 13 autologous BMT or T cell depleted allogeneic BMT recipients undergoing treatment for relapsed leukemia or lymphoma. Enhanced immunological responsiveness was demonstrated in the laboratory with a 5- to 40-fold increase in circulating CD56 bright CD16⁺ CD3⁻ NK cells. Moreover, this low dose regimen of IL-2 resulted in augmented in vitro killing of the NK targets K562.

25 When Soiffer et al. (55) updated the outcome of 29 allogeneic BMT patients who received low dose IL-2, they found superior survival for these patients (70%) compared to histological controls (30%, $p=0.41$).

-27-

Lauria et al. (56) treated 11 patients with high grade NHL at a median of 42 days after ABMT with IL-2 at a dose of 2×10^6 IU/m² qod for two weeks and then 3×10^6 IU/m² twice a week for a year. Phenotypic analysis showed a persistent and significant ($p=0.001$) increase in the proportion and absolute
5 number of total lymphocytes and especially of both CD16 and CD56 NK cells after 6 months of therapy. None of the patients progressed with a median follow-up of twenty-two months (range 10-42 months) after starting therapy. In addition, two patients with residual disease after ABMT, one in the liver and second in the lymph nodes, obtained a complete response after 7 and 10 months of IL-2 therapy.

10 Vey et al. (57) treated 25 patients with refractory or relapsed HD (11 patients) and NHL (14 patients) with low dose IL-2. 48% of the patients had resistant disease at transplant and 84% achieved CR after ABMT. IL-2 was started at a mean of 54 days after transplant and consisted of a first cycle of 5 days followed by 4 cycles of 2 days every other week. Patients received a mean of 160
15 $\times 10^6$ IU/m² of IL-2. After a five year follow-up, the probability of survival and DFS is 72% (HD 73% and NHL 70%) and 45% (HD 36% and NHL 48%).

A group at the Fred Hutchinson Cancer Research Center (FHCRC) has recently found that low dose IL-2 therapy was well-tolerated in the outpatient setting, and that remissions in patients treated with low dose IL-2 tended to be
20 longer than without IL-2 treatment. IL-2 therapy was associated with an increase in the number of certain populations of immune cells, including CD8+ CD69+ cells; CD16+ CD8+ cells; CD16+ CD69+ cells; CD16+ CD56+ cells; CD16+ CD122+ cells; CD16+ Dr+ cells; and CD8+ CD56+ cells. There was also an increase in the expression of lytic activity against the tumor targets K562 and
25 Daudi, with a median of 5.9-fold and 6.5-fold increase, respectively. Relapses, when they occurred, occurred at a median of 17.8 months after transplant, and

-28-

therefor remissions were reported to be characteristically longer than what was historically seen in transplant recipients without IL-2 therapy.

Given the encouraging data gathered from single therapy studies with IL-2 on ABMT transplant recipients, it seemed reasonable to combine IL-2 therapy with Rituximab® post transplant, given that Rituximab's biological activity appears to be mediated through ADCC and complement-mediated lytic activity. Thus, a Phase I trial has been initiated in collaboration with the FHCRC to evaluate the safety and potential efficacy of a combined therapeutic regimen.

A separate Phase II study is also being performed to evaluate the efficacy and the incidence of HACA formation in patients receiving low-dose IL-2 and Rituxan®. A specific objective of this study is to assess whether ADCC is enhanced by in vivo exposure to IL-2 and whether ADCC activity correlates with clinical response. Inclusion criteria for patients are histologically confirmed stage II-IV low-grade, follicular B-cell or mantle cell lymphoma. Mantle cell lymphoma, for the purposes of this clinical study, is defined as CD5+, CD23- (if available) and/or bcl-1+ by immunohistochemistry. Patients who did not respond to or have relapsed following their first treatment with a standard therapy, i.e., chemotherapy, radiotherapy, ABMT and/or immunotherapy, are eligible.

Rituximab® plus GM-CSF for the Treatment of Relapsed Low Grade or Follicular B-Cell Lymphoma

Two separate Phase II trials have also been initiated to test the efficacy of combined treatment with Rituximab® and GM-CSF. One study involves 40 patients with relapsed low grade B-cell lymphoma, and comprises administering Rituximab® at 375 mg/m² weekly X 4 (d. 1, 8, 15, 22) and GM-CSF (Leukine, Immunex) at 250 mcg sc three times weekly for 8 weeks, starting one hour before the first dose of Rituximab®. This study will be used to evaluate the clinical efficacy (overall response rate (ORR), overall complete response rate, time to

-29-

progression and failure-free survival) of the combined therapeutic regimen, to characterize the safety (qualitative, quantitative, duration and reversibility of adverse events) of the combined therapy, and to determine the effects of the combined therapy on relevant lymphocyte subsets and cytokines. The second
5 study plans to also monitor immunologic parameters to assess the mechanism of killing (complement C3 and C4, CH50, flow cytometry for CD3, CD4, CD8, CD16, CD19 and CD56 and ADCC assay).

Rituximab® plus Gamma-Interferon

Gamma-interferon may also be useful in combined therapy with
10 Rituximab® for treating patients with low-grade or higher-grade lymphomas. It is has recently been found that gamma-interferon upregulates CD20 expression on multiple myeloma (MM) patient plasma cells, patient B-cells, as well as on normal donor B-cells (Treon et al., Lugano, 1999). In fact, Treon and colleagues have shown that gamma-interferon augments binding of these cells to Rituximab®.
15 Induction of CD20 expression on plasma cells occurred in a dose dependent manner, with upregulation seen with as little as 1 U/ml of interferon gamma. A plateau occurred at 100 U/ml at 48 hours. Thus, gamma-interferon may also be beneficial when administered in combination with Rituximab®.

INTERMEDIATE-GRADE AND HIGH-GRADE NHL

Single-Agent Studies

In a study conducted in Europe and Australia, alternative dosing schedules were evaluated in 54 relapsed or refractory intermediate- or high-grade NHL patients (34). Rituximab® was infused at 375 mg/m² weekly for 8 doses or at 375 mg/m² once followed by 500 mg/m² weekly for 7 doses. The ORR was 31%; (CR
25 9%, PR 22%) no significant difference between the dosing regimens was observed. Patients with diffuse large-cell lymphoma (N = 30) had an ORR of 37% and those with mantle-cell lymphoma (N = 12) had an ORR of 33%.

-30-

Combination of Rituximab® and CHOP Chemotherapy

In another study, 31 patients with intermediate- or high-grade NHL (19 females, 12 males, median age 49) received Rituximab® on Day 1 of each of six 21 -day cycles of CHOP (35). Of 30 evaluable patients, there were 19 CR (63%) and 10 PR (33%), for an ORR of 96%. This regimen was considered well tolerated and may result in higher response rates than with Rituximab® or CHOP alone.

The NCI Division of Cancer Treatment and Diagnosis is collaborating with IDEC Pharmaceuticals Corporation to explore Rituximab® treatment in other indications. A Phase II trial of CHOP versus CHOP and Rituximab® is being conducted by ECOG, CALGB, and SWOG in older patients (> 60 years) with mixed, diffuse large cell, and immunoblastic large cell histology NHL (N = 630 planned). This study includes a secondary randomization to maintenance with Rituximab® versus non-maintenance.

A Phase III trial of Rituximab® and CHOP in 40 patients with previously untreated mantle-cell lymphoma is also ongoing at the Dana Farber Institute. Rituximab® is administered on Day 1 and CHOP is given on Days 1 - 3 every 21 days for 6 cycles. Accrual for this study has been completed. A Phase II trial of CHOP followed by Rituximab® in newly diagnosed follicular lymphoma conducted by SWOG has also been completed. Results of these two trials are being analyzed.

A Phase II trial of CHOP and Rituximab® versus CHOP alone in HIV-related NHL conducted by the AIDS Malignancy Consortium is ongoing; 120 patients are planned.

Rituximab® after Myeloablative Therapy Relapse

Rituximab® has shown promising early results in patients with relapsed intermediate-grade NHL after high-dose therapy with autologous PBSC support.

-31-

Six of seven patients responded (1 CR and 5 PR) and one patient had stable disease; therapy was well tolerated (36).

SAFETY EXPERIENCE

Adverse events and clinical laboratory data from 315 patients in the five
5 single-agent U.S. studies were combined to provide a safety profile of
Rituximab® in patients with low-grade or follicular NHL. The majority of
adverse events were infusion-related and occurred with decreasing frequency after
the first infusion. The most common infusion-related events were fever (49%),
chills (32%), nausea (18%), fatigue (16%), headache (14%), angioedema (13%),
10 pruritus (10%), and occasionally, hypotension (10%) and bronchospasm (8%).
During the treatment period (up to 30 days following the last dose), 10% of
patients experienced Grade 3 or 4 adverse events, which were primarily infusion-
related or hematologic. Thrombocytopenia ($< 50,000$ platelets/mm³) occurred in
1.3% of patients, neutropenia (< 1000 /mm³) occurred in 1.9%, and anemia (< 8
15 gm/dL) occurred in 1.0%. Although Rituximab® induced B-cell depletion in 70%
- 80% of patients, abnormally decreased serum immunoglobulins were observed
in a minority of patients and the incidence of infection did not appear to be
increased.

Hypotension requiring interruption of the Rituximab® infusion occurred
20 in 10% of patients and was Grade 3 or 4 in 1 %. Angioedema was reported in
13% of patients and was considered serious in one patient. Bronchospasm
occurred in 8% of patients; 2% were treated with bronchodilators. A single report
of bronchiolitis obliterans was noted. Most patients experienced no further
infusion-related toxicities by the second and subsequent infusions. The
25 percentage of patients reporting adverse events upon retreatment was similar to
that reported following the first course(14).

-32-

Four patients developed arrhythmias during Rituximab® infusion. One of the four discontinued treatment because of ventricular tachycardia and supraventricular tachycardias. The other three patients experienced trigeminy (N = 1) and irregular pulse (N = 2) and did not require discontinuation of therapy.

5 Angina was reported during infusion and myocardial infarction occurred four days postinfusion in one subject with a prior history of myocardial infarction.

The overall incidence of adverse events and Grade 3 and 4 adverse events was higher in patients with bulky disease than in patients with non-bulky disease. The incidence of dizziness, neutropenia, thrombocytopenia, myalgia, anemia, and chest pain was higher in patients with lesions > 10 cm. The incidence of Grade

10 3 or 4 neutropenia, anemia, hypotension, and dyspnea was also higher in patients with bulky disease compared with patients with lesions < 10 cm (19).

Since FDA approval of Rituximab® for treatment of relapsed or refractory low-grade or follicular NHL in 1997, an estimated 17,000 patients have been treated. In May, 1998, descriptions of eight post-marketing reports of severe

15 infusion-related adverse events associated with the use of Rituximab® that resulted in fatal outcomes were summarized. In seven of the eight fatalities, severe symptoms occurred during the first Rituximab® infusion. The cause of death was not reported or remains unknown for two of the eight cases. Severe

20 respiratory events, including hypoxia, pulmonary infiltrates, or adult respiratory distress syndrome contributed to six of the eight reported deaths. One patient had a pretreatment lymphocyte count of 600,000/mm³; another, a creatinine of 8; a third, a respiratory rate of 40; and a fourth, pancytopenia. Patients with a high tumor burden or with a high number of circulating malignant cells may be at

25 higher risk and these patients should be monitored closely throughout each infusion.

-33-

Most of the adverse events recently described were previously observed in Rituximab® clinical studies. One notable exception is an infusion-related syndrome associated with rapid tumor lysis, that was reported in six patients with high numbers of circulating tumor cells (37,38). This syndrome was characterized by fever, rigors, bronchospasm with associated hypoxemia, a rapid decline in peripheral lymphocytes, laboratory evidence of tumor destruction, and transient, severe thrombocytopenia. These patients had diagnoses of B-prolymphocytic leukemia (N = 2), chronic lymphocytic leukemia (N = 2), mantle-cell lymphoma (N = 1), or transformed NHL (N = 1); all had elevated circulating lymphocytes, bulky adenopathy, and organomegaly. Although five of these six patients required hospitalization, symptoms resolved and subsequent Rituximab® treatments were well tolerated; the last patient refused further therapy and died of progressive disease two weeks later.

In a separate report of seven patients with CLL and one patient with mantle-cell lymphoma, tumor lysis syndrome was observed after the first Rituximab® infusion in those patients with lymphocyte counts $> 10 \times 10^9/L$ (39).

RADIOIMMUNOTHERAPY WITH ^{90}Y TTTRIUM-LABELED ANTI-CD20 ANTIBODY IN COMBINATION WITH RITUXIMAB®

Another therapeutic approach to NHL under evaluation is a radiolabeled anti-CD20 antibody (IDEC-Y2B8) in combination with Rituximab®. IDEC-Y2B8 (^{90}Y -ibritumomab tiuxetan) is a murine IgG₁ kappa anti-CD20 antibody conjugated to ^{90}Y via a chelator, MX-DTPA, which is covalently bound to the antibody. Rituximab® (250 mg/m²) is administered prior to IDEC-Y2B8 to deplete peripheral B lymphocytes and improve biodistribution of the radiolabeled antibody.

In a recently reported Phase I/II study (40-42), patients with low-grade NHL (N = 34), intermediate-grade NHL (N = 14), or mantle-cell lymphoma (N

-34-

= 3) were treated with IDEC-Y2B8. The median age was 60, 71% were male, and 96% were Caucasian. Of 51 patients with relapsed or refractory NHL, 34 (67%) responded to single doses of 0.2, 0.3, or 0.4 mCi/kg of IDEC-Y2B8. The ORR was 82% (28/34) for patients with low-grade or follicular NHL and was 43% (6/14) for patients with intermediate-grade lymphoma. No patients with mantle-cell disease responded.

A Phase III randomized study comparing IDEC-Y2B8 with Rituximab® (375 mg/m² weekly times 4) for treatment of low-grade follicular or transformed NHL patients is ongoing. Another Phase III trial is also being conducted in patients with relapsed NHL who are refractory to Rituximab®.

SUMMARY

In the absence of curative therapy for NHL, the objective of treatment is to achieve control of the disease for a meaningful duration and provide relief of tumor-related symptoms without undue toxicity. Treatment with Rituximab® is a brief, 22-day outpatient therapy with limited adverse events in most patients. In clinical studies, 50% of evaluable relapsed or chemotherapy refractory low-grade or follicular NHL patients achieved complete or partial responses. These responses were durable without maintenance therapy; the median TTP for responders was 13.2 months and the median DR was 11.6 months in the pivotal study.

Rituximab® is approved as a safe and effective treatment for patients with relapsed low-grade or follicular B-cell NHL. It has significant clinical activity, a novel mechanism of action, and compares favorably with alternative therapies in response rate and response duration. Completion of ongoing studies will verify the role of alternative Rituximab® regimens and Rituximab® in the treatment of other CD20+ B-lymphocyte malignancies.

References

1. Press O, Appelbaum F, Ledbetter J, Martin P, Zarling J, Kidd P, Thomas E. Monoclonal antibody 1F5 (anti-CD20) serotherapy of human B-cell lymphomas. *Blood* 1987; 69:584-591.
- 5 2. Dillman R. Antibodies as cytotoxic therapy. *Journal of Clinical Oncology* 1994; 12:1497-1515.
- 3 Grossbard M, Press O, Appelbaum F, Bernstein I, Nadler L. Monoclonal antibody-based therapies of leukemia and lymphoma. *Blood* 1992; 80:863-878.
- 10 4. Reff M, Camer K, Chambers K, Chinn P, Leonard J, Raab R, Newman R, Hanna N, Anderson D. Depletion of B cells in vivo by a chimeric mouse human monoclonal antibody to CD20. *Blood* 1994; 83:435-445.
5. Demidem A, Lam T, Alas S, Hariharan K, Hanna N, Bonavida B. Chimeric anti-CD20 (IDEC-C2B8) monoclonal antibody sensitizes a B cell lymphoma cell line to cell killing by cytotoxic drugs. *Cancer Biotherapy & Radiopharmaceuticals* 1997; 12:177-186.
- 15 6. Maloney D, Liles T, Czerwinski D, Waldichuk C, Rosenberg J, Grillo-López A, Levy R. Phase I clinical trial using escalating single-dose infusion of chimeric anti-CD20 monoclonal antibody (IDEC-C2B8) in patients with recurrent B-cell lymphoma. *Blood* 1994; 84:2457-2466.
- 20 7. Maloney D, Grillo-López A, White C, Bodkin D, Schilder R, Neidhart J, Janakiraman N, Foon K, Liles T-M, Dallaire B, Wey K, Royston I, Davis T, Levy R. IDEC-C2B8 (Rituximab®) anti-CD20 monoclonal antibody therapy in patients with relapsed low-grade non-Hodgkin's lymphoma. *Blood* 1997; 90: 2188-2195.
- 25 8. McLaughlin P, Grillo-López A, Link B, Levy R, Czuczman M, Williams M, Heyman M, Bence-Bruckler I, White C, Cabanillas F, Jain V, Ho A,

-36-

- Lister J, Wey K, Shen D, Dallaire B. Rituximab® chimeric anti-CD20 monoclonal antibody therapy for relapsed indolent lymphoma: half of patients respond to a 4-dose treatment program. *Journal of Clinical Oncology* 1998; 16:2825-2833.
- 5 9. McLaughlin P, Grillo-López A, Maloney D, Link B, Levy R, Czuczman M, Cabanillas F, Dallaire B, White C. Efficacy controls in long-term follow-up of patients treated with rituximab for relapsed or refractory, low-grade or follicular NHL. *Blood* 1998; 92:414a-415a.
- 10 10. Janakiraman N, McLaughlin P, White C, Maloney D, Shen D, Grillo-López A. Rituximab: Correlation between effector cells and clinical activity in NHL. *Blood* 1998; 92 (10 Suppl 1):337a.
- 15 11. Berinstein N, Grillo-López A, White C, Bence-Bruckler I, Maloney D, Czuczman M, Green D, Rosenberg J, McLaughlin P, Shen D. Association of serum Rituximab (IDEC-C2B8) concentration and anti-tumor response in the treatment of recurrent low-grade or follicular non-Hodgkin's lymphoma. *Annals of Oncology* 1998; 9:995-1001.
- 20 12. Tobinai K, Kobayashi Y, Narabayashi M, Ogura M, Kagami Y, Morishima Y, Ohtsu T, Igarashi T, Sasaki Y, Kinoshita T, Murate T. Feasibility and pharmacokinetic study of a chimeric anti-CD20 monoclonal antibody (IDEC-C2B8, rituximab) in relapsed B-cell lymphoma. *Annals of Oncology* 1998; 9:527-534.
- 25 13. Piro L, White C, Grillo-López A, Janakiraman N, Saven A, Beck T, Varns C, Shuey S, Czuczman M, Lynch J, Kolitz J, Jain V. Extended Rituxan (anti-CD20 monoclonal antibody) therapy for relapsed or refractory low-grade or follicular non-Hodgkin's lymphoma. 1999; Submitted
14. Davis T, White C, Grillo-López A, Velasquez W, Link B, Maloney D, Dillman R, Williams M, Mohrbacher A, Weaver R, Dowden S, Levy R.

- Rituximab: First report of a Phase II (PII) trial in NHL patients (pts) with bulky disease. *Blood* 1998; 92 (10 Suppl 1):414a.
15. Byrd J, White C, Thomas S, Veldsquez W, Rosenberg J, Grillo-López A. Rituximab therapy in previously treated Waldenstrom's Macroglobulinemia: Preliminary evidence of activity. *Blood* 1998; 92 (IO Suppl 1): 106(a).
16. O'Brien S, Freireich E, Andreeff M, Lemer S, Keating M. Phase I/III Study of Rituxan in chronic lymphocytic leukemia (CLL). *Blood* 1998; 92:105a, #431.
- 10 17. Venugopal P, Sivaraman S, Huang X, Chopra H, O'Brein T, Jajeh A, Preisler H. Upregulation of CD20 expression in chronic lymphocytic leukemia (CLL) cells by *in vitro* exposure to cytokines. *Blood* 1998; 10:247a.
18. Flinn I, O'Donnell P, Noga S, Vogelsang G, Grever M, Goodrich A, Abrams R, Marcellus D, Miller C, Jones R, Ambinder R. *In vivo* purging and adjuvant immunotherapy with Rituximab PBSC transplant for NHL. *Blood* 1998; 92:648a, #2673.
- 15 19. Davis T, Levy R, White C, Czuczman M, McLaughlin P, Link B, Varns C, Weaver R, Grillo-López A. Rituximab: Phase II (PII) retreatment (ReRx) study in patients (pts) with low-grade or follicular (LG/F) NHL. *Blood* 1998; 92 (10 Suppl 1):414a.
- 20 20. Davis T, Czerwinski D, Levy R. Therapy of B cell lymphoma with anti-CD20 antibodies can result in the loss of CD20 antigen expression. *Clinical Cancer Research* 1999; 5: In press.
- 25 21. Czuczman M, Grillo-López A, White C, Saleh M, Gordon L, LoBuglio F, Jonas C, Klippenstein D, Dallaire B, Varns C. Treatment of patients with low-grade B-cell lymphoma with the combination of chimeric anti-CD20

-38-

- monoclonal antibody and CHOP chemotherapy. *Journal of Clinical Oncology* 1999; 17:268-276.
22. White C, Czuczman M, Grillo-López A, White C, Saleh M, Gordon L, LoBuglio A, Jonas C, Alkuzweny B, Dowen S. Rituximab/CHOP chemoimmunotherapy in patients (pts) with low grade lymphoma (LG/F NHL): Progression free survival (PFS) after three years (median) follow-up. *Proceedings of ASCO* 1999, In press.
23. Wadler S, Schwartz E. Principles in the biomodulation of cytotoxic drugs by interferons. *Seminars in Oncology* 1992; 19:45-48.
- 10 24. Nakamura K, Kubo A, Hosokawa S, Nagaike K, Hashimoto S. Effect of alpha-interferon on anti-alpha-fetoprotein-monoclonal-antibody targeting of hepatoma. *Oncology* 1993; 50:35-40.
25. Greiner J, Guadagni F, Noguchi P, Pestka S, Colcher D, Fisher P, Schlom J. Recombinant interferon enhances monoclonal antibody-targeting of carcinoma lesions in vivo. *Science* 1987; 235:895-898.
- 15 26. Murray J, Zukiwski A, Mujoo K, Rosenblum M. Recombinant alpha-interferon enhances tumor targeting of an antimelanoma monoclonal antibody in vivo. *Journal of Biological Response Modifiers* 1990; 9:556-563.
- 20 27. Yokota S, Hara H, Luo Y, Seon B. Synergistic potentiation of in vivo antitumor activity of anti-human T-leukemia immunotoxins by recombinant alpha-interferon and daunorubicin. *Cancer Research* 1990; 50:32-37.
28. Grillo-López A, Dallaire B, Shen C, Varns C, McClure A, Caralli V. Treatment options for patients with relapsed low-grade or follicular lymphoma: The role of IDEC-C2B8. *Antibody Imunoconjugates and Radiopharmaceuticals* 1995; 8:60.
- 25

29. Davis T, Maloney D, White C, Grillo-López A, Williams M, Weiner G, Sklenar T, Levy R. Combination immunotherapy of low grade or follicular (LG/F) non-Hodgkin's lymphoma (NHL) with Rituximab and alpha interferon: Interim analysis. *Proceedings of the American Society of Clinical Oncology* 1998; 17:11a.
30. Smalley R, Andersen J, Hawkins M, Bhide V, O'Connell M, Oken M, Borden E. Interferon alfa combined with cytotoxic chemotherapy for patients with non-Hodgkin's lymphoma. *New England Journal of Medicine* 1992; 327: 1336-1341.
31. Hagenbeek A, Carde P, Meerwaldt JH, Somers R, Thomas J, De Bock R, Raemaekers JM, van Hoof A, De Wolf-Peeters C, van Glabbeke M. Maintenance of remission with human recombinant interferon alfa-2a in patients with stages In and IV low-grade malignant non-Hodgkin's lymphoma. European Organization for Research and Treatment of Cancer Lymphoma Cooperative Group. *Journal of Clinical Oncology* 1998; 16:41-47.
32. Solal-Céligny P, Lepage E, Brousse N, Tendler C, Brice P, Haioun C, Gabarre J, Pignon B, Tertian G, Bouabdallah R, Rossi J-F, Doyen C, Coiffier B. Doxorubicin-containing regimen with or without interferon alfa-2b for advanced follicular lymphomas: Final analysis of survival and toxicity in the groupe d'etude des lymphomes folliculaires 86 trial. *Journal of Clinical Oncology* 1998; 16:2332-2338.
33. van der Kolk L, Grillo-López A, Gerritsen W, Jonkhoff A, Baars J, van Oers M. Chimeric anti-CD20 monoclonal antibody (rituximab) plus G-CSF in relapsed B-cell lymphoma: A phase I/II clinical trial. *Blood* 1998; 92:241b, #4037.

-40-

34. Coiffier B, Haioun C, Ketterer N, Engert A, Tilly H, Ma D, Johnson P, Lister A, Feuring-Buske M, Radford JA, Capdeville R, Diehl V, Reyes F. Rituximab (anti-CD20 monoclonal antibody) for the treatment of patients with relapsing or refractory aggressive lymphoma: a multicenter phase H
5 study. *Blood* 1998; 92:1927-1932.
35. Link B, Grossbard M, Fisher R, Czuczman M, Gilman P, Lowe A, Vose J. Phase II pilot study of the safety and efficacy of rituximab in combination with CHOP chemotherapy in patients with previously
10 untreated- or high-grade NHL. *Proceedings of the American Society of Clinical Oncology* 1998; 17:3a.
36. Tsai, D, Moore H, Porter D, Vaughn D, Luger S, Loh R, Schuster S, Stadtmauer E. Progressive intermediate grade non-Hodgkin's lymphoma after high dose therapy and autologous peripheral stem cell transplantation (PSCT) has a high response rate to Rituximab. *Blood* 1998; 92:415a,
15 #1713.
37. Byrd J, Waselenko J, Maneatis T, Murphy T, Weickum R, Ward F, White C. Rituximab therapy in hematologic malignancy patients with circulating blood tumor cells: Association with increased infusion-related side effects and rapid tumor lysis. *Blood* 1998; 92 (IO Suppl 1): 106a.
- 20 38. Jensen M, Winkler U, Manzke O, Diehl V, Engert A. Rapid tumor lysis in a patient with B-cell chronic lymphocytic leukemia and lymphocytosis treated with an anti-CD20 monoclonal antibody (IDEC- C2B8, rituximab). *Annals of Hematology* 1998; 77:89-91.
- 25 39. Winkler U, Jensen M, Manzke O, Tesch H, Bohlen H, Diehl V, Engert A. Severe side effects in patients with B-cell chronic lymphocytic leukemia (CLL) and lymphocytosis treated with the monoclonal antibody Rituximab. *Blood* 1998; 92:285b, #4228.

-41-

40. Witzig T, White C, Wiseman G, Gordon L, Emmanouilides C, Raubitschek A, Janakiraman N, Gutheil J, Spies S, Silverman D, Parker E, Grillo-López A. Phase I/II trial of IDEC-Y2B8 radioimmunotherapy for treatment of relapsed or refractory CD20 positive B-cell non-Hodgkin's lymphoma. *Journal of Clinical Oncology* 1999; Submitted.
41. Wiseman G, White C, Witzig T, Gordon L, Emmanouilides C, Raubitschek A, Janakiraman N, Spies S, Silverman D, Gutheil J, Schilder R, Parker E, Rosenberg J, Grillo-López A. IDEC-Y2B8 radioimmunotherapy: Baseline bone marrow involvement and platelet count are better predictors of hematologic toxicity than dosimetry. *Blood* 1998; 92:417a.
42. Witzig T, White C, Wiseman G, Gordon L, Emmanouilides C, Raubitschek A, Janakiraman N, Spies S, Silverman D, Gutheil J, Schilder R, Ding E, Shen D, Grillo-López A. IDEC-Y2B8 Radioimmunotherapy: Responses in patients with splenomegaly. *Blood* 1998; 92:417(a).
43. Witherspoon RP, Lum LG, Storb R. Immunologic reconstitution after bone marrow grafting. *Semin Hematol* 21:2, 1984.
44. Anderson, KC et al. Hematological engraftment and immune reconstitution posttransplant with anti-B1 purged autologous bone marrow. *Blood* 69:597, 1987.
45. Lum LG. Kinetics of immune reconstitution after human marrow transplantation. *Blood* 69:369, 1987.
46. Azogui O., Gluckman E., Fradelizi, D., Inhibition of IL-2 production after human allogeneic bone marrow transplantation. *J. Immunol.* 131:1205, 1983
47. Welte, K. et al, Defective Interleukin-2 production in patients after bone marrow transplantation and in vitro restoration of defective T lymphocyte proliferation by highly purified Interleukin. *Blood* 64:380, 1984.

-42-

48. Cayeau, S. et al., T-cell ontogeny after bone marrow transplantation: failure to synthesize Interleukin-2 (IL-2) and lack of CD2- and CD3-mediated proliferation by both CDE4+ and CD8+ cells even in the presence of exogenous IL-2. *Blood* 74:2270, 1989.
- 5 49. Bosley, A. et al., Interleukin-2 as consolidative immuotherapy against minimal residual disease. *Nouv Rev Fr Hematol* 32:13, 1990.
50. Caligiuri, M.A. et al, Extended continuous infusion low-dose recombinant Interleukin-2 in advanced cancer. Prolonged immunomodulation without significant toxicity. *J Clin Oncol* 9:2110, 1991.
- 10 51. Caligiuri, M.A. et al, Selective immune modulation of NK cells following prolonged infusions of low dose recombinant IL-2. *J Clin Invest* 91:123, 1993.
52. Caligiuri, M.A., Low-dose reocmbinant Interleukin-2 therapy: rationale and potential clinical applications. *SEM in Oncol* 20:3, 1993.
- 15 53. Klarnet, J.P. et al, Antigen-driven T cell clones can proliferate in vivo, eradicate disseminated leukemia and provide specific immunologic memory. *J Immunol.* 138:4012, 1987.
54. Soiffer, R.J. et al, Clinical and immunologic effects of prolonged infusion of low-doe recombinant Interleukin-2 after autologous and T cell-depleted
20 allogeneic bone marrow transplantation. *Blood* 79:517, 1992.
55. Soiffer, R.J. et al, Effect of low-dose Interleukin-2 on disease relapse after T-cell depleted allogeneic bone marrow transplantation. *Blood* 84:964, 1994.
56. Lauria, F. et al, Immunologic and clinical modifications following low-
25 dose subcutaneous administration of rIL-2 in non-Hodgkin's lymphoma patients after autologous bone marrow transplantation. *BMT* 18:79, 1996.

-43-

57. Vey, N. et al, A pilot study of autologous bone marrow transplantation followed by recombinant Interleukin-2 in malignant lymphomas. *Leukemia & Lymphoma* 21:107, 1996.
 58. Venugopal, P. et al, Upregulation of CD20 expression in CLL cells by cytokines. Submitted to ASH Meeting, December 1998.
- 5

WHAT IS CLAIMED:

1. A method for treating relapsed B-cell lymphoma comprising administering to a patient having relapsed the B-cell lymphoma a therapeutically effective amount of an anti-CD20 antibody.

5 2. The method of Claim 1, wherein said patient was treated previously with an anti-CD20 antibody.

3. The method of Claim 1, wherein said patient previously underwent a bone marrow or stem cell transplantation.

4. The method of Claim 1, wherein said patient previously underwent
10 radiotherapy.

5. The method of Claim 1, wherein said patient previously underwent chemotherapy for said B-cell lymphoma.

6. The method of Claim 5, wherein said chemotherapy is selected from the group consisting of CHOP, ICE, Mitozantrone, Cytarabine, DVP, ATRA,
15 Idarubicin, hoelzer chemotherapy regime, La La chemotherapy regime, ABVD, CEOP, 2-CdA, FLAG & IDA with or without subsequent G-CSF treatment), VAD, M & P, C-Weekly, ABCM, MOPP and DHAP.

7. A method for treating a subject having B-cell lymphoma, which subject has not exhibited appreciable tumor remission or regression after
20 administration of a chimeric anti-CD20 antibody, comprising administering to said patient a radiolabeled anti-CD20 antibody.

8. The method of Claim 7, wherein said radiolabeled anti-CD20 antibody is administered from about one week to about two years after said administration of said chimeric anti-CD20 antibody.

25 9. The method of Claim 8, wherein said radiolabeled anti-CD20 antibody is administered from about one week to about nine months after said administration of said chimeric anti-CD20 antibody.

-45-

10. The method of Claim 1, wherein said anti-CD20 antibody is a chimeric anti-CD20 antibody.

11. The method of Claim 10, wherein said chimeric antibody is C2B8 (Rituximab®).

5 12. A method for treating B-cell lymphoma comprising administering a synergistic therapeutic combination comprising at least one anti-CD20 antibody and at least one cytokine, wherein the therapeutic effect is better than the additive effects of either therapy administered alone.

10 13. The method of Claim 12, wherein said at least one cytokine is selected from the group consisting of alpha interferon, gamma interferon, IL-2, GM-CSF and G-CSF.

14. The method of Claim 13, wherein said anti-CD20 antibody and said alpha interferon, gamma interferon, IL-2, GM-CSF or G-CSF is administered sequentially, in either order, or in combination.

15 15. The method of Claim 12, wherein said anti-CD20 antibody is a chimeric antibody.

16. The method of Claim 15, wherein said chimeric anti-CD20 antibody is C2B8 (Rituximab®).

20 17. A method for treating B-cell lymphoma comprising administering to a patient a therapeutically effective amount of anti-CD20 antibody before, during or subsequent to a chemotherapeutic regimen.

25 18. The method of Claim 17, wherein said chemotherapy regimen is selected from the group consisting of CHOP, ICE, Mitozantrone, Cytarabine, DVP, ATRA, Idarubicin, hoelzer chemotherapy regime, La La chemotherapy regime, ABVD, CEOP, 2-CdA, FLAG & IDA with or without subsequent G-CSF treatment), VAD, M & P, C-Weekly, ABCM, MOPP and DHAP.

-46-

19. The method of Claim 17, wherein said anti-CD20 antibody is a chimeric antibody.

20. The method of Claim 19, wherein said chimeric antibody is C2B8 (Rituximab®).

5 21. A method for treating B-cell lymphoma comprising administering to a patient a therapeutically effective amount of an anti-CD20 antibody before, during or subsequent to a bone marrow or stem cell transplant.

22. The method of Claim 21, wherein said anti-CD20 antibody is a chimeric anti-CD20 antibody.

10 23. The method of Claim 22, wherein said chimeric anti-CD20 antibody is C2B8 (Rituximab®).

24. A method of reducing residual CD20+ tumor cells in bone marrow or stem cells before or after myeloablative therapy by administering to a patient an anti-CD20 antibody.

15 25. The method of Claim 24, wherein said anti-CD20 antibody is a chimeric anti-CD20 antibody.

26. The method of Claim 25, wherein said chimeric anti-CD20 antibody is C2B8 (Rituximab®).

20 27. The method of Claim 1, wherein said B-cell lymphoma is selected from the group consisting of low grade/follicular non-Hodgkin's lymphoma (NHL), small lymphocytic (SL) NHL, intermediate grade/follicular NHL, intermediate grade diffuse NHL, chronic lymphocytic leukemia (CLL), high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL, mantle cell lymphoma, AIDS-related
25 lymphoma and Waldenstrom's Macroglobulinemia.

28. The method of Claim 12, wherein said B-cell lymphoma is selected from the group consisting of low grade/ follicular non-Hodgkin's lymphoma

(NHL), small lymphocytic (SL) NHL, intermediate grade/ follicular NHL, intermediate grade diffuse NHL, chronic lymphocytic leukemia (CLL), high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL, mantle cell lymphoma, AIDS-related lymphoma and Waldenstrom's Macroglobulinemia.

29. The method of Claim 17, wherein said B-cell lymphoma is selected from the group consisting of low grade/follicular non-Hodgkin's lymphoma (NHL), small lymphocytic (SL) NHL, intermediate grade/follicular NHL, intermediate grade diffuse NHL, chronic lymphocytic leukemia (CLL), high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL, mantle cell lymphoma, AIDS-related lymphoma and Waldenstrom's Macroglobulinemia.

30. The method of Claim 21, wherein said B-cell lymphoma is selected from the group consisting of low grade/follicular non-Hodgkin's lymphoma (NHL), small lymphocytic (SL) NHL, intermediate grade/follicular NHL, intermediate grade diffuse NHL, chronic lymphocytic leukemia (CLL), high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL, mantle cell lymphoma, AIDS-related lymphoma and Waldenstrom's Macroglobulinemia.

31. The method of Claim 24, wherein said B-cell lymphoma is selected from the group consisting of low grade/follicular non-Hodgkin's lymphoma (NHL), small lymphocytic (SL) NHL, intermediate grade/follicular NHL, intermediate grade diffuse NHL, chronic lymphocytic leukemia (CLL), high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL, mantle cell lymphoma, AIDS-related lymphoma and Waldenstrom's Macroglobulinemia.

